

Discovery of 10,059 new three-dimensional periodic orbits of general three-body problem

Xiaoming Li¹ and Shijun Liao^{2,3*}

¹School of Mechanics and Construction Engineering, Jinan University, Guangzhou 510632, China

²School of Ocean and Civil Engineering, Shanghai Jiao Tong University, Shanghai 200240, China

³School of Physics and Astronomy, Shanghai Jiao Tong University, Shanghai 200240, China

*Corresponding author: sjliao@sjtu.edu.cn

Abstract *A very few three-dimensional (3D) periodic orbits of general three-body problem (with three finite masses) have been discovered since Newton mentioned it in 1680s. Using a high-accuracy numerical strategy we discovered 10,059 three-dimensional periodic orbits of the three-body problem in the cases of $m_1 = m_2 = 1$ and $m_3 = 0.1n$ where $1 \leq n \leq 20$ is an integer, among which 1,996 (about 20%) are linearly stable. Note that our approach is valid for arbitrary mass m_3 so that in theory we can gain an arbitrarily large amount of 3D periodic orbits of the three-body problem. In the case of three equal masses, we discovered twenty-one 3D “choerographical” periodic orbits whose three bodies move periodically in a single closed orbit. It is very interesting that, in the case of two equal masses, we discovered 273 three-dimensional periodic orbits with the two bodies ($m_1 = m_2 = 1$) moving along a single closed orbit and the third ($m_3 \neq 1$) along a different one: we name them “piano-trio” orbits, like a trio for two violins and one piano. To the best of our knowledge, all of these 3D periodic orbits have never been reported, indicating the novelty of this work. The large amount of these new 3D periodic orbits are helpful for us to have better understandings about chaotic properties of the famous three-body problem, which “are, so to say, the only opening through which we can try to penetrate in a place which, up to now, was supposed to be inaccessible”, as pointed out by Poincaré, the founder of chaos theory.*

Keywords general three-body problem, three-dimensional periodic orbits

1. INTRODUCTION

The famous three-body problem can be traced back to Newton [1] in 1680s, which focuses on the motion of three bodies with finite masses that attract to each other by gravitational force under Newton’s second law. It attracted many famous mathematicians and physicists such as Euler [2] and Lagrange [3] who found a very few of the earliest planar periodic orbits of the three-body problem. The three-body problem is three-dimensional (3D) in essence, but in the three hundred years after Newton mentioned it, a very few periodic orbits of the general three-body problem were found almost in the two-dimensional (2D) case that is certainly much simpler than the three-dimensional one. It was Poincaré [4] who revealed its mathematical complexity and difficulty in 1890: the first integrals for the motion of the three-body system do not exist so that its closed-form solution is impossible in general, and besides its trajectories have the sensitivity dependence on initial conditions that laid the foundation of modern chaos theory. This well explains why in the 300 years only three families of planar periodic orbits of three-body system were found by Euler [2] in 1767 and Lagrange [3] in 1772, until 1970s when the Broucke-Hadjidemetriou-Henon family of planar periodic orbits [5–7] were discovered by means of computer. In 1993 the so-called Figure-8 planar periodic orbit was numerically discovered by Moore [8] and subsequently rediscovered and rigorously proven by Chenciner and Montgomery [9], which is called “choreographic” orbit: the three bodies move periodically in a single closed orbit. In 2013, Šuvakov and Dmitrašinović [10] made a breakthrough to numerically discover 13 new distinct planar periodic orbits by means of modern computer using Eulerian collinear initial configuration that has since been widely adopted to search for planar periodic orbits [11–14].

Why is it so difficult to find periodic orbits of three-body system even in 2013 when there exist supercomputers with very high performance? It was Lorenz [15] who revealed the reason in 2006: despite the sensitivity dependence on initial condition [4], a chaotic system has also the sensitivity dependence on numerical algorithm so that chaotic trajectory is quickly polluted by numerical noise. Thus, since three-body problem is essentially chaotic according to Poincaré [4], its trajectory should be sensitive to numerical noises that are unavoidable for traditional numerical algorithms such as Runge-Kutta’s method (in double precision) that is widely used to gain trajectories of three-body system. In 2009 Liao [16] proposed the so-called “clean numerical method” (CNS) [17–21] whose numerical noises can be reduced to such a low level that accurate/convergent chaotic trajectories can be gained in a finite but long enough time interval $t \in [0, T_c]$, where the so-called “predictable critical time” T_c is determined by the level of background numerical noise.




Figure 1: The initial configuration of the 3D three-body system. The initial positions of the three bodies are located at $\mathbf{r}_1 = (-1, 0, 0)$, $\mathbf{r}_2 = (1, 0, 0)$ and $\mathbf{r}_3 = (0, 0, z_0)$ with the corresponding initial velocities $\mathbf{v}_1 = (v_x, v_y, v_z)$, $\mathbf{v}_2 = (v_x, v_y, -v_z)$ and $\mathbf{v}_3 = (-2v_x/m_3, -2v_y/m_3, 0)$.

We emphasize here that accurate chaotic trajectory is very important for three-body problem, since it is the key reason why the number of discovered new families of planar periodic orbits of general three-body problem increases to several orders of magnitude: in 2017 Li and Liao [22] discovered more than 600 new families of planar periodic orbits of three-body problem with three equal masses by means of computer using a numerical strategy based on CNS, then in 2018 Li, Jing and Liao [23] further discovered 1,223 new families of planar periodic orbits of three-body problem with two equal masses in a similar way, in 2021 Li et al. [24] successfully obtained 135,445 new planar periodic orbits (including 13,315 stable ones) with arbitrarily unequal masses, and in 2022 Liao et al. [25] proposed an effective roadmap to numerically gain planar periodic orbits of three-body systems with arbitrary masses by means of CNS and machine learning. In 2024 Hristov et al. [26] found 24,582 equal-mass periodic orbits of free-fall three-body problem by means of high-accuracy algorithm. Finally, there are no any obstacles for us to gain masses of new *planar* periodic orbits of three-body system with arbitrary masses by means of high-accuracy numerical algorithms such as CNS.

However, compared to the planar periodic orbits mentioned above, the three-dimensional (3D) periodic orbits of three-body problem are far limited. Two types of periodic orbits were mainly studied. The first is the spatial isosceles three-body problem, in which two bodies have equal masses and the third body oscillates along a straight line, while the three bodies always form an isosceles triangle. In the so-called restricted three-body problem that contains one massless body, this configuration is known as the Sitnikov problem [27], where two massive bodies move along an elliptical orbit in a plane, and the massless body oscillates in a straight line perpendicular to the plane. This class of 3D periodic orbits has been extended to the general three-body problem, where all three bodies have finite masses [28–30]. The second types were gained using the continuation methods to generate 3D periodic orbits from periodic orbits of restricted three-body problem [31, 32]. It is a great pity that a very few 3D periodic orbits of general three-body problem have been found, to the best of our knowledge. Thus, it has great scientific meanings to discover a masses of 3D periodic orbits of general three-body problem with finite masses.

2. INITIAL CONFIGURATION AND NUMERICAL STRATEGY

The motions of the general three-body system (with finite masses) are governed by the dimensionless equations

$$(1) \quad \frac{d^2 \mathbf{r}_i}{dt^2} = - \sum_{j=1, j \neq i}^3 m_j \frac{(\mathbf{r}_i - \mathbf{r}_j)}{|\mathbf{r}_i - \mathbf{r}_j|^3}, \quad i = 1, 2, 3,$$

where t denotes the time, \mathbf{r}_i and m_i are the position vector and mass of the i -th body, respectively. Let $\mathbf{X}(t) = (\mathbf{r}_1(t), \mathbf{r}_2(t), \mathbf{r}_3(t), \dot{\mathbf{r}}_1(t), \dot{\mathbf{r}}_2(t), \dot{\mathbf{r}}_3(t))$ denote the state space of the three-body system at the time t , where $\dot{\mathbf{r}}_i(t)$ represents the velocity of the i -th body. To evaluate the proximity of the system to its initial state $\mathbf{X}(0)$, we define the so-called “return proximity function”

$$(2) \quad \delta(t) = |\mathbf{X}(t) - \mathbf{X}(0)|.$$

A periodic orbit is determined by $\delta(T) = 0$, where T is its period.




Figure 2: 3D periodic orbits of general three-body problem: (a) $O_2(1.2)$, (b) $O_8(0.6)$, (c) $O_3(1.0)$, (d) $O_4(1.0)$, (e) $O_6(1.0)$, (f) $O_6(1.2)$. Blue: Body-1; Red: Body-2; Green: Body-3. Filled circle: initial positions of the three bodies.

As shown in Fig. 1, we use an initial configuration for the 3D three-body system in this paper. The initial positions of the three bodies are located at $\mathbf{r}_1 = (-1, 0, 0)$, $\mathbf{r}_2 = (1, 0, 0)$ and $\mathbf{r}_3 = (0, 0, z_0)$, the initial velocities of Body-1 and Body-2 are $\mathbf{v}_1 = (v_x, v_y, v_z)$ and $\mathbf{v}_2 = (v_x, v_y, -v_z)$, respectively, where z_0, v_x, v_y, v_z are unknown. For the sake of simplicity, let us consider the case of zero momentum, i.e. $m_1\mathbf{v}_1 + m_2\mathbf{v}_2 + m_3\mathbf{v}_3 = 0$, which gives the initial velocity $\mathbf{v}_3 = (-(m_1 + m_2)v_x/m_3, -(m_1 + m_2)v_y/m_3, 0)$ of Body-3. Note that in the above-mentioned configuration the initial condition of the 3D three-body system is determined by the four unknown variables v_x, v_y, v_z , and z_0 .

Our numerical strategy is briefly described as follows. First of all, we use the grid search method to search for possible candidates of 3D periodic orbits of the three-body system in the case of $v_z = 0$ within $z_0 \in (0, 1]$, $v_x \in (0, 1]$ and $v_y \in (0, 1]$: 1000 isometric points are used in each dimension, i.e. for a chosen m_3 totally $1000 \times 1000 \times 1000 = 10^9$ cases are considered as initial condition to integrate the governing equations (1) using the DOP853 solver [33]. One among these initial conditions is regarded as a candidate of 3D periodic orbit if its corresponding return proximity function (2) is less than 10^{-1} under a value of period T . Secondly, the values of (z_0, v_x, v_y, v_z) and T of each candidate are constantly modified/corrected by means of the numerical approach (similar to [23]) based on the Newton-Raphson method [34] and Clean Numerical Simulation (CNS) [16–21]: a 3D periodic orbit is discovered when the return proximity function $\delta(T)$ defined by (2) is less than 10^{-10} . The linear stability of periodic orbits is evaluated by Floquet theory [35]. To compute the monodromy matrix, we integrate the variational equations along periodic orbit [36]. When all eigenvalues of the monodromy matrix lie on the unit circle, the periodic orbit is linearly stable.

3. DISCOVERY OF 10,059 THREE-DIMENSIONAL PERIODIC ORBITS

Note that m_3 is a variable although $m_1 = m_2 = 1$ is fixed. Without loss of generality, we consider the case $m_3 = 0.1n$ where $1 \leq n \leq 20$ is an integer, respectively, thus totally 2×10^{10} cases are done for the the grid search method. Using a national supercomputer, we discovered 10,059 three-dimensional periodic orbits of the general three-body system. It is found that 1,996 among them are linearly stable, approximately 20% of the total. Their initial condition, period T and stability are given on the website <https://github.com/sjtu-liao/three-body>. These 3D periodic orbits are named $O_n(m_3)$, where n corresponds to the n -th orbit when ordered by its value of period T (from small to large) for a fixed mass m_3 (when $m_1 = m_2 = 1$). For example, $O_1(1)$ denotes the discovered 3D periodic orbit with the shortest period T




Figure 3: 3D choreographic periodic orbits of three-body problem in case of $m_1 = m_2 = m_3 = 1$: (a) $O_{62}(1.0)$, (b) $O_{64}(1.0)$, (c) $O_{231}(1.0)$, (d) $O_{524}(1.0)$. Three bodies move along a single closed orbit (blue). Filled circles: the initial positions of the three bodies. Blue: Body-1; Red: Body-2; Green: Body-3.

when $m_3 = 1$. Some 3D periodic orbits are shown in Fig. 2 and their corresponding initial conditions (z_0, v_x, v_y, v_z) and periods T are given in Table S1 in Supplementary Information. To the best of our knowledge, all of these 3D periodic orbits have *never* been reported and thus are novel. It should be emphasized that we can choose an *arbitrary* value of m_3 , such as $m_3 = 0.615$ or $m_3 = 0.872$, to obtain 3D periodic orbits in the similarly way. So, in theory, we can gain an *arbitrarily* large amount of 3D periodic orbits of three-body system using the same initial configuration as that mentioned in this paper.

In the case of three equal masses, i.e. $m_1 = m_2 = m_3 = 1$, we discover 1,504 three-dimensional periodic orbits. Among them, there are 21 “choreographic” orbits, say, the three bodies move periodically in a single closed orbit. Their initial conditions, periods and stability are given in Table S2 in Supplementary Information. Different from the famous planar choreographic orbit Figure-8 [8, 9], these twenty-one choreographic orbits are *three-dimensional*, as shown in Fig. 3. To the best of our knowledge, such kind of 3D choreographic periodic orbits of the general three-body system have *never* been reported.

Besides, it is very interesting that, when $m_1 = m_2 = 1$ and $m_3 \neq 1$, we found 273 three-dimensional periodic orbits in the 19 cases of $m_3 = 0.1n$ ($1 \leq n \leq 20$ but $n \neq 10$ is an integer), where the two bodies with equal mass ($m_1 = m_2 = 1$) move along a single closed orbit but the third ($m_3 \neq 1$) moves along a different closed orbit. We name them “piano-trio orbit”, like a trio for two violins and one piano. Their initial conditions, periods and stability are given on the website <https://github.com/sjtu-liao/three-body>. Four orbits of them are as shown in Fig. 4, where Body-1 ($m_1 = 1$) and Body-2 ($m_2 = 1$) follow a single closed orbit (in red) but Body-3 ($m_3 \neq 1$) traces a distinct path (in green), and their initial conditions, periods and stability are given in Table S3 in Supplementary Information. To the best of our knowledge, such kind of 3D periodic orbits of general three-body system have *never* been reported.

4. CONCLUDING REMARKS AND DISCUSSION

In this paper, by means of a numerical strategy based on the grid search method, Newton-Raphson method [34] and clean numerical method (CNS) [16–21], we successfully discovered 10,059 three-dimensional periodic orbits of the three-body problem in the cases of $m_1 = m_2 = 1$ and $m_3 = 0.1n$ ($1 \leq n \leq 20$), among which 1,996 (about 20%) are




Figure 4: 3D “piano-trio orbits” of three-body problem with two bodies ($m_1 = m_2 = 1$) moving along a single closed orbit (red) but the third ($m_3 \neq 1$) along a distinct orbit (green): (a) $O_6(0.6)$, (b) $O_{26}(1.1)$, (c) $O_{48}(0.5)$, (d) $O_{267}(0.9)$. Filled circles: the initial positions of the three bodies. Blue: Body-1 ($m_1 = 1$) ; Red: Body-2 ($m_2 = 1$) ; Green: Body-3 ($m_3 \neq 1$).

linearly stable. It should be emphasized that our numerical approach is valid for *arbitrary* mass m_3 (such as $m_3 = 0.615$ or $m_3 = 0.872$) so that in theory we can gain an *arbitrarily* large amount of 3D periodic orbits of the three-body problem in a similar way (we will do it and update the dataset on the website <https://github.com/sjtu-liao/three-body>). Certainly, these 3D periodic orbits should be helpful for us to have better understandings about the chaotic properties of the famous three-body problems, since they “are, so to say, the only opening through which we can try to penetrate in a place which, up to now, was supposed to be inaccessible”, as pointed out by Poincaré, the founder of chaos theory.

It should be emphasized that, in the case of three equal masses ($m_1 = m_2 = m_3 = 1$), we discovered twenty-one 3D “choerographical” periodic orbits whose three bodies move periodically in a single closed orbit. To the best of our knowledge, these 3D “choerographical” periodic orbits have *never* been reported, which reveals its general existence for three-body problem and indicates the novelty of this work. Besides, in the case of two equal masses ($m_1 = m_2 = 1$ and $m_3 \neq 1$), we discovered 273 three-dimensional periodic orbits with two bodies ($m_1 = m_2 = 1$) moving along a single closed orbit and the third $m_3 \neq 1$ along a different one: we name them “piano-trio” orbits, like a trio for two violins and one piano. To the best of our knowledge, these “piano-trio” orbits have *never* been reported, indicating the novelty of this work.

As illustrated by Liao et al. [25], one can gain arbitrarily accurate planar periodic orbit of three-body problem by means of CNS. Similarly, each of our discovered 10,059 three-dimensional periodic orbits reported in this paper could be in an arbitrary accuracy. For example, the high accuracy (in 70 significant digits) initial condition and the period T of the linearly stable periodic orbit $O_3(1.0)$ are listed in Table S4 in Supplementary Information. In physics, the minimum spatial distance is a Planck length $l_p = 1.616252 \times 10^{-35}$ meter and the maximum distance is the diameter of the observable universe (as the characteristic length) $d_u \approx 9.3 \times 10^{10}$ light year $\approx 8.4 \times 10^{26}$ meter, so that the minimum dimensionless spatial length that has physical meaning is $l_p/d_u \approx 1.9 \times 10^{-62}$. Note that our 3D periodic orbit $O_3(1.0)$ listed in Table S4 in Supplementary Information is accurate in the 70 significant digits, which thus can be regarded, from the physical viewpoint, as accurate as a closed-form solution.

Many further investigations should be done in future. It is true that in this paper we only consider the case with two equal masses $m_1 = m_2 = 1$. However, using the known 3D periodic orbits obtained in this paper as a starting point, it is straightforward to gain new 3D periodic orbits of general three-body problem with three *unequal* masses by means of the continuation method [24]. Note that, for planar periodic orbits of three-body problem, their topological identification can be classified by braid groups [8] or shape-space sphere [10, 37]. However, for a 3D periodic orbit of general three-body problem, it is an open question how to identify its topological classification [8].

ACKNOWLEDGMENT

The calculations were performed on the Tianhe-2 Supercomputer, National Supercomputer Center in Guangzhou, China. This work is partly supported by National Natural Science Foundation of China (No. 12002132) and State Key Laboratory of Ocean Engineering.

REFERENCES

- [1] I. Newton, *Philosophiæ naturalis principia mathematica* (Mathematical principles of natural philosophy), London: Royal Society Press, 1687.
- [2] L. Euler, *De motu rectilineo trium corporum se mutuo attrahentium*, *Novi commentarii academiae scientiarum Petropolitanae* (1767) 144–151.
- [3] J.-L. Lagrange, *Essai sur le probleme des trois corps*, *Prix de l'académie royale des Sciences de paris* 9 (1772) 292.
- [4] J. H. Poincaré, *Sur le problème des trois corps et les équations de la dynamique*. Divergence des séries de M. Lindstedt, *Acta Math.* 13 (1890) 1–270.
- [5] R. Broucke, H. Lass, *A note on relative motion in the general three-body problem*, *Celestial mechanics* 8 (1) (1973) 5–10.
- [6] J. D. Hadjidemetriou, T. Christides, *Families of periodic orbits in the planar three-body problem*, *Celestial mechanics* 12 (2) (1975) 175–187.
- [7] M. Hénon, *A family of periodic solutions of the planar three-body problem, and their stability*, *Celestial mechanics* 13 (3) (1976) 267–285.
- [8] C. Moore, *Braids in classical dynamics*, *Phys. Rev. Lett.* 70 (1993) 3675–3679.
- [9] A. Chenciner, R. Montgomery, *A remarkable periodic solution of the three-body problem in the case of equal masses*, *Annals of Mathematics* 152 (3) (2000) 881–901.
- [10] M. Šuvakov, V. Dmitrašinović, *Three classes of newtonian three-body planar periodic orbits*, *Phys. Rev. Lett.* 110 (2013) 114301.
- [11] P. P. Iasko, V. V. Orlov, *Search for periodic orbits in the general three-body problem*, *Astronomy Reports* 58 (11) (2014) 869–879. doi:10.1134/S1063772914110080. URL <https://doi.org/10.1134/S1063772914110080>
- [12] A. Hudomal, *New periodic solutions to the three-body problem and gravitational waves*, M.S. Thesis, University of Belgrade, Serbia (2015).
- [13] M. Šuvakov, M. Shibayama, *Three topologically nontrivial choreographic motions of three bodies*, *Celestial Mechanics and Dynamical Astronomy* 124 (2) (2016) 155–162.
- [14] D. Rose, *Geometric phase and periodic orbits of the equal-mass, planar three-body problem with vanishing angular momentum*, Ph.D. Thesis, University of Sydney, Australia (2016).
- [15] E. N. Lorenz, *Computational periodicity as observed in a simple system*, *Tellus A: Dynamic Meteorology and Oceanography* 58A (2006) 549 – 557.
- [16] S. Liao, *On the reliability of computed chaotic solutions of non-linear differential equations*, *Tellus A* 61 (4) (2009) 550–564.
- [17] S. Liao, P. Wang, *On the mathematically reliable long-term simulation of chaotic solutions of lorenz equation in the interval [0,10000]*, *Sci. China - Phys. Mech. Astron.* 57 (2014) 330 – 335.
- [18] X. Li, S. Liao, *Clean numerical simulation: a new strategy to obtain reliable solutions of chaotic dynamic systems*, *Applied Mathematics and Mechanics* 39 (11) (2018) 1529–1546.
- [19] T. Hu, S. Liao, *On the risks of using double precision in numerical simulations of spatio-temporal chaos*, *Journal of Computational Physics* 418 (2020) 109629.
- [20] S. Qin, S. Liao, *Influence of numerical noises on computer-generated simulation of spatio-temporal chaos*, *Chaos Solitons Fractals* 136 (2020) 109790.
- [21] S. Liao, *Clean Numerical Simulation*, CRC Press, 2023.
- [22] X. Li, S. Liao, *More than six hundred new families of newtonian periodic planar collisionless three-body orbits*, *SCIENCE CHINA Physics, Mechanics & Astronomy* 60 (12) (2017) 129511.
- [23] X. Li, Y. Jing, S. Liao, *Over a thousand new periodic orbits of a planar three-body system with unequal masses*, *Publications of the Astronomical Society of Japan* 70 (4) (2018) 64.
- [24] X. Li, X. Li, S. Liao, *One family of 13315 stable periodic orbits of non-hierarchical unequal-mass triple systems*, *Science China Physics, Mechanics & Astronomy* 64 (1) (2021) 219511.
- [25] S. Liao, X. Li, Y. Yang, *Three-body problem: From newton to supercomputer plus machine learning*, *New Astronomy* 96 (2022) 101850.
- [26] I. Hristov, R. Hristova, V. Dmitrašinović, K. Tanikawa, *Three-body periodic collisionless equal-mass free-fall orbits revisited*, *Celestial Mechanics and Dynamical Astronomy* 136 (1) (2024) 7.
- [27] K. Sitnikov, *The Existence of Oscillatory Motions in the Three-Body Problem*, *Soviet Physics Doklady* 5 (1961) 647.

- [28] M. Corbera, J. Llibre, Families of periodic orbits for the spatial isosceles 3-body problem, *SIAM Journal on Mathematical Analysis* 35 (5) (2004) 1311–1346. doi:10.1137/S0036141002407880.
URL <http://dx.doi.org/10.1137/S0036141002407880>
- [29] D. Yan, R. Liu, X. Hu, W. Mao, T. Ouyang, New phenomena in the spatial isosceles three-body problem with unequal masses, *International Journal of Bifurcation and Chaos* 25 (12) (2015) 1550169.
- [30] O. M. Perdomo, A bifurcation in the family of periodic orbits for the spatial isosceles 3 body problem, *Qualitative Theory of Dynamical Systems* 17 (2) (2017) 411–428. doi:10.1007/s12346-017-0244-1.
URL <http://dx.doi.org/10.1007/s12346-017-0244-1>
- [31] K. Katopodis, Continuation of periodic orbits: Three-dimensional circular restricted to the general three-body problem, *Celestial Mechanics* 19 (1) (1979) 43–51. doi:10.1007/bf01230173.
URL <http://dx.doi.org/10.1007/BF01230173>
- [32] V. V. Markellos, The three-dimensional general three-body problem: Determination of periodic orbits, *Celestial Mechanics* 21 (3) (1980) 291–309. doi:10.1007/bf01230227.
URL <http://dx.doi.org/10.1007/BF01230227>
- [33] E. Hairer, G. Wanner, S. P. Norsett, *Solving Ordinary Differential Equations I: Non-stiff Problems*, Springer-Verlag Berlin Heidelberg, 1993.
- [34] A. Abad, R. Barrio, A. Dena, Computing periodic orbits with arbitrary precision, *Phys. Rev. E* 84 (2011) 016701.
- [35] J. Guckenheimer, P. Holmes, *Nonlinear Oscillations, Dynamical Systems, and Bifurcations of Vector Fields*, Springer New York, 1983. doi:10.1007/978-1-4612-1140-2.
URL <http://dx.doi.org/10.1007/978-1-4612-1140-2>
- [36] K. Richter, G. Tanner, D. Wintgen, Classical mechanics of two-electron atoms, *Physical Review A* 48 (6) (1993) 4182.
- [37] R. Montgomery, The n-body problem, the braid group, and action-minimizing periodic solutions, *Nonlinearity* 11 (2) (1998) 363.

Supplementary Information

Table S1: The initial conditions, the periods T and the stability of some 3D periodic orbits in the case of the initial position $\mathbf{r}_1 = (-1, 0, 0)$, $\mathbf{r}_2 = (1, 0, 0)$ and $\mathbf{r}_3 = (0, 0, z_0)$ with the initial velocities $\mathbf{v}_1 = (v_x, v_y, v_z)$, $\mathbf{v}_2 = (v_x, v_y, -v_z)$ and $\mathbf{v}_3 = (-2v_x/m_3, -2v_y/m_3, 0)$. Stability of periodic orbit: S (linearly stable), U (linearly unstable).

Orbit $O_n(m_3)$	z_0	v_x	v_y	v_z	T	stability
$O_2(1.2)$	1.0220057827E+00	-2.7260000746E-01	-4.3209371195E-01	6.2947340717E-01	6.9057763983E+00	U
$O_8(0.6)$	7.2379454056E-01	2.1745309714E-01	-2.4023509732E-01	4.8072135301E-01	8.2513531298E+00	U
$O_3(1.0)$	4.7687826428E-01	4.0213691007E-01	1.8035695129E-01	2.1044512814E-01	6.8316220363E+00	S
$O_4(1.0)$	1.0656465072E-01	4.1396073100E-01	4.7046903637E-02	4.4299118421E-02	7.2303054580E+00	S
$O_6(1.0)$	2.5738110869E-01	3.0241724398E-01	5.6056948094E-01	5.2896109217E-03	1.3654015842E+01	U
$O_6(1.2)$	4.4254976548E-01	6.3300867799E-01	1.7485764338E-01	-2.6069495427E-01	8.2359759431E+00	U

Table S2: The initial conditions, the periods T and the stability of 21 choreographic 3D periodic orbits having three bodies moving along the same path in the case of the initial position $\mathbf{r}_1 = (-1, 0, 0)$, $\mathbf{r}_2 = (1, 0, 0)$ and $\mathbf{r}_3 = (0, 0, z_0)$ with the initial velocities $\mathbf{v}_1 = (v_x, v_y, v_z)$, $\mathbf{v}_2 = (v_x, v_y, -v_z)$ and $\mathbf{v}_3 = (-2v_x/m_3, -2v_y/m_3, 0)$. Stability of periodic orbit: S (linearly stable), U (linearly unstable).

Orbit $O_n(m_3)$	z_0	v_x	v_y	v_z	T	stability
$O_{62}(1.0)$	1.4254560021E-01	3.4973952580E-01	6.0245969580E-01	5.6258558122E-03	4.4413258151E+01	U
$O_{64}(1.0)$	2.4089819317E-01	2.9736744570E-01	5.5127998288E-01	6.4720700746E-03	4.5160913797E+01	U
$O_{231}(1.0)$	1.3449345804E-01	3.3746407711E-01	5.3495391505E-01	1.7586983000E-03	8.2638396171E+01	S
$O_{264}(1.0)$	1.2527262314E-01	3.3812559500E-01	5.3478083136E-01	1.8426277977E-03	8.8847804260E+01	S
$O_{468}(1.0)$	1.9882559981E-01	3.5789268529E-01	5.3413197456E-01	-5.3389716378E-04	1.1603279244E+02	U
$O_{524}(1.0)$	9.2921053426E-02	3.4199333600E-01	5.3399733168E-01	1.6826415762E-03	1.2041512314E+02	S
$O_{574}(1.0)$	2.4895906866E-01	3.3871794052E-01	5.5917249111E-01	-2.5336443257E-03	1.2535891398E+02	S
$O_{609}(1.0)$	7.9082023338E-02	2.8653274947E-01	5.3295320774E-01	1.3330149482E-03	1.2885423891E+02	U
$O_{617}(1.0)$	2.5059548392E-01	3.1883837350E-01	5.5145840743E-01	7.2285919945E-04	1.2927664287E+02	U
$O_{623}(1.0)$	2.4884379653E-01	2.9277777970E-01	5.5413846299E-01	6.9945423951E-03	1.2975160287E+02	U
$O_{735}(1.0)$	1.4523478543E-01	3.3263557496E-01	5.2616069788E-01	-3.1374383107E-03	1.4041031143E+02	U
$O_{793}(1.0)$	1.5066856881E-01	3.3509372155E-01	5.3528992956E-01	1.2102376067E-03	1.4629315855E+02	U
$O_{941}(1.0)$	1.3957142176E-01	3.3672076170E-01	5.3505271584E-01	1.6586446900E-03	1.5893652668E+02	S
$O_{1034}(1.0)$	7.9952100998E-02	3.9991383731E-01	5.2892786472E-01	2.6785166522E-03	1.6684804313E+02	U
$O_{1062}(1.0)$	2.5387857578E-01	3.0855862740E-01	5.5506896654E-01	3.4527925512E-03	1.6887732550E+02	S
$O_{1114}(1.0)$	1.9719159455E-01	3.4208132539E-01	5.3782184023E-01	1.4597658877E-03	1.7279868625E+02	S
$O_{1172}(1.0)$	6.3233041014E-02	3.4473818853E-01	5.3330581391E-01	1.2154707447E-03	1.7726783502E+02	S
$O_{1265}(1.0)$	6.1061814942E-02	3.4489838389E-01	5.3326623797E-01	1.1774765759E-03	1.8358838190E+02	S
$O_{1414}(1.0)$	4.9792913732E-02	2.6873050476E-01	5.2603515916E-01	3.2150647009E-04	1.9557410885E+02	S
$O_{1488}(1.0)$	1.5788544119E-01	3.3561402713E-01	5.3228206363E-01	-2.2004697254E-03	2.0197023210E+02	S
$O_{1497}(1.0)$	1.1019540397E-01	3.3940414208E-01	5.3491778263E-01	2.0614879729E-03	2.0302355527E+02	S

Table S3: The initial conditions, the periods T and the stability of some 3D periodic orbits having two bodies moving along the same path in the case of the initial position $\mathbf{r}_1 = (-1, 0, 0)$, $\mathbf{r}_2 = (1, 0, 0)$ and $\mathbf{r}_3 = (0, 0, z_0)$ with the initial velocities $\mathbf{v}_1 = (v_x, v_y, v_z)$, $\mathbf{v}_2 = (v_x, v_y, -v_z)$ and $\mathbf{v}_3 = (-2v_x/m_3, -2v_y/m_3, 0)$. Stability of periodic orbits: S (linearly stable), U (linearly unstable).

Orbit $O_n(m_3)$	z_0	v_x	v_y	v_z	T	stability
$O_6(0.6)$	6.1460435884E-01	1.0381048232E-01	7.2494633230E-02	5.8966905104E-02	7.9743488400E+00	U
$O_{26}(1.1)$	6.1717279647E-01	5.0563910986E-01	2.8923907397E-01	6.3478431464E-03	2.7381502412E+01	U
$O_{48}(0.5)$	2.5356269746E-01	3.0249330624E-01	7.6131511629E-02	-3.9679960521E-03	9.0118410122E+01	S
$O_{267}(0.9)$	5.7288742602E-01	3.2057036195E-01	2.3346447921E-01	3.1450924543E-03	8.6966037898E+01	S

Table S4: The high accuracy (in 70 significant digits) initial condition and the period T of the linearly stable periodic orbit $O_3(1.0)$ in the case of the initial position $\mathbf{r}_1 = (-1, 0, 0)$, $\mathbf{r}_2 = (1, 0, 0)$ and $\mathbf{r}_3 = (0, 0, z_0)$ with the initial velocities $\mathbf{v}_1 = (v_x, v_y, v_z)$, $\mathbf{v}_2 = (v_x, v_y, -v_z)$, $\mathbf{v}_3 = (-2v_x/m_3, -2v_y/m_3, 0)$ and $m_3 = 1.0$.

Orbit $O_3(1.0)$	
z_0	4.768782642803115460616110933545915887541644925960260369253641303299294E-1
v_x	4.021369100747237493832253390929016257978052115111599603699588130212840E-1
v_y	1.803569512862586641317199478891917026060004921964179954624832790547685E-1
v_z	2.104451281378731221497174690764795110575771159319962594676582115390083E-1
T	6.831622036284445040766025062746064085737914286107653347931784678966507